

On the Solution of the Number-Projected Hartree-Fock-Bogoliubov Equations

J.A. Sheikh⁽¹⁾, E. Lopes⁽¹⁾ and P. Ring⁽¹⁾

⁽¹⁾*Physik-Department, Technische Universität München, D-85747 Garching bei München,
Germany*

Abstract

The numerical solution of the recently formulated number-projected Hartree-Fock-Bogoliubov equations is studied in an exactly soluble cranked-deformed shell model Hamiltonian. It is found that the solution of these number-projected equations involve similar numerical effort as that of bare HFB. We consider that this is a significant progress in the mean-field studies of the quantum many-body systems. The results of the projected calculations are shown to be in almost complete agreement with the exact solutions of the model Hamiltonian. The phase transition obtained in the HFB theory as a function of the rotational frequency is shown to be smeared out with the projection.

PACS numbers : 21.60.Cs, 21.10.Hw, 21.10.Ky, 27.50.+e

Typeset using REVTeX

The mean-field models with effective forces have been quite successful in describing the gross features of the quantum many-body systems. Although, the mean-field approaches are appropriate for systems with a very large number of particles, they have also been quite useful to describe the properties of finite quantum systems, for instance the atomic nucleus. The ground-state properties of atomic nuclei have been well described using the Hartree-Fock (HF) and Hartree-Fock-Bogoliubov (HFB) mean-field approaches with various effective interactions [1]. However, the mean-field application to a finite system suffers from a fundamental problem that it leads to sharp phase transitions. The phase transition is an artefact of the mean-field approach and is not observed in the experimental data. The phase transition obtained is due to the neglect of the quantal fluctuations, which become quite strong for finite systems.

There are various methods in the literature to consider the quantal fluctuations on the mean-field solution for the finite system. One very powerful method is through the restoration of the broken-symmetries by employing the projection methods [1]. In the present work, we shall consider the restoration of the gauge-symmetry associated with the particle-number. It is known in the HFB studies that one often obtains a phase transition from the superfluid paired-phase to the normal unpaired-phase. This phase transition is due to the fluctuations in the particle-number, the HFB wavefunction does not have a well defined particle-number. In most of the analysis, the particle-number fluctuations are treated in an approximate way by employing the Lipkin-Nogami prescription [2-4]. However, it has been shown that this approach also breaks down at high-rotational frequencies and as a matter of fact violates the variational principle [5].

The exact particle number-projection can be performed by using the gradient methods [1]. But this approach is numerically quite involved and has been applied to only separable interactions with restricted model spaces [6,7]. There has been an unresolved issue whether HFB like equations can be obtained with the projected- energy functional. This problem has been recently solved [8] and it has been shown that it is possible to obtain the HFB equations from an arbitrary real energy functional which is completely expressible in terms of

the density-matrix and the pairing-tensor. The projected-energy functional can be expressed in terms of the density-matrix and the pairing-tensor and one obtains the HFB equations with modified expressions for the pair-gap and the Hartree-Fock potential. The expressions for these quantities acquire a relatively simple form for the case of particle number-projection [8].

To check the applicability of the number projected-HFB (PHFB) formalism, detailed numerical analysis is carried out in a simple cranked-deformed shell model Hamiltonian [9]. Although, this model cannot be used directly to study the experimental data, but it contains all the basic ingredients of a more realistic model. The advantage in this model is that it can be solved exactly and it is possible to check the accuracy of an approximate method. We consider that it is quite instructive to test the number-projection method in a cranking model as the Coriolis forces destroy the pair-correlations and the results become quite sensitive to the treatment of the pairing-interaction. As we shall demonstrate, the present projection method reproduces almost exactly the results of the shell model calculations for all the cases studied.

The model Hamiltonian consists of a cranked deformed one-body term, h' and a scalar two-body delta-interaction [9]. The one-body term is the familiar cranked-Nilsson mean-field potential which takes into account of the long-range part of the nucleon-nucleon interaction. The residual short-range interaction is specified by the delta-interaction. The deformed shell model Hamiltonian employed is given by

$$H = h'_{def} + V_2, \quad (1)$$

$$H = h_{def} - \omega J_x + V_2, \quad (2)$$

where,

$$h_{def} = -4\kappa\sqrt{\frac{4\pi}{5}} \sum_m \langle jm|Y_{20}|jm \rangle c_{jm}^\dagger c_{jm}, \quad (3)$$

and

$$V_2 = \frac{1}{2} \sum_{LM} E_L A_{LM}^\dagger A_{LM}, \quad (4)$$

with $A_{LM}^\dagger = (c_j^\dagger c_j^\dagger)_{LM}$ and $A_{LM} = (A_{LM}^\dagger)^\dagger$. For the antisymmetric-normalized two-body matrix-element (E_J), we use the delta-interaction which for a single j-shell is given by [10]

$$E_L = -G \frac{(2j+1)^2}{2(2L+1)} \begin{bmatrix} j & j & L \\ \frac{1}{2} & -\frac{1}{2} & 0 \end{bmatrix}^2, \quad (5)$$

where the symbol $[\]$ denotes the Clebsch-Gordon coefficient. We use $G = g \int R_{nl}^4 r^2 dr$ as our energy unit and the deformation energy κ is related to the deformation parameter β . For the case of $h_{11/2}$ shell, $\kappa=2.4$ approximately corresponds to $\beta = 0.23$.

It has been shown in ref. [8] that the variation of the number projected-energy functional results in the HFB matrix equation

$$\mathcal{H}' \begin{pmatrix} U \\ V \end{pmatrix} = E'_i \begin{pmatrix} U \\ V \end{pmatrix}, \quad (6)$$

where

$$\mathcal{H}' = \begin{pmatrix} \varepsilon'_{n_1 n_2} + \Gamma_{n_1 n_2} + \Lambda_{n_1 n_2} - \lambda \delta_{n_1 n_2} & \Delta_{n_1 n_2} \\ -\Delta_{n_1 n_2}^* & -\varepsilon'^*_{n_1 n_2} - \Gamma_{n_1 n_2}^* - \Lambda_{n_1 n_2}^* + \lambda \delta_{n_1 n_2} \end{pmatrix}. \quad (7)$$

The number-projected expressions for $\varepsilon'_{n_1 n_2}, \Gamma_{n_1 n_2}, \Lambda_{n_1 n_2}$ and $\Delta_{n_1 n_2}$ are given by

$$\begin{aligned} \varepsilon' &= \frac{1}{2} \int d\phi y(\phi) \left(Y(\phi) \text{Tr}[e' \rho(\phi)] + [1 - 2ie^{-i\phi} \sin \phi \rho(\phi)] e' C(\phi) \right) \\ &+ h.c. \end{aligned} \quad (8)$$

$$\begin{aligned} \Gamma &= \frac{1}{2} \int d\phi y(\phi) \left(Y(\phi) \frac{1}{2} \text{Tr}[\Gamma(\phi) \rho(\phi)] + \frac{1}{2} [1 - 2ie^{-i\phi} \sin \phi \rho(\phi)] \Gamma(\phi) C(\phi) \right) \\ &+ h.c. \end{aligned} \quad (9)$$

$$\begin{aligned} \Lambda &= -\frac{1}{2} \int d\phi y(\phi) \left(Y(\phi) \frac{1}{2} \text{Tr}[\Delta(\phi) \bar{\kappa}^*(\phi)] - 2ie^{-i\phi} \sin \phi C(\phi) \Delta(\phi) \bar{\kappa}^* \right) \\ &+ h.c. \end{aligned} \quad (10)$$

$$\Delta = \frac{1}{2} \int d\phi y(\phi) e^{-2i\phi} C(\phi) \Delta(\phi) - (..)^\dagger, \quad (11)$$

with

$$\Gamma_{n_1 n_3}(\phi) = \sum_{n_2 n_4} \bar{v}_{n_1 n_2 n_3 n_4} \rho_{n_4 n_2}(\phi), \quad (12)$$

$$\Delta_{n_1 n_2}(\phi) = \frac{1}{2} \sum_{n_3 n_4} \bar{v}_{n_1 n_2 n_3 n_4} \kappa_{n_3 n_4}(\phi), \quad (13)$$

$$\bar{\Delta}_{n_3 n_4}^*(\phi) = \frac{1}{2} \sum_{n_1 n_2} \bar{\kappa}_{n_1 n_2}^*(\phi) \bar{v}_{n_1 n_2 n_3 n_4}, \quad (14)$$

$$\rho(\phi) = C(\phi) \rho, \quad (15)$$

$$\kappa(\phi) = C(\phi) \kappa = \kappa C^T(\phi), \quad (16)$$

$$\bar{\kappa}(\phi) = e^{2i\phi} \kappa C^*(\phi) = e^{2i\phi} C^\dagger(\phi) \kappa, \quad (17)$$

$$C(\phi) = e^{2i\phi} \left(1 + \rho(e^{2i\phi} - 1) \right)^{-1}, \quad (18)$$

$$x(\phi) = \frac{1}{2\pi} \frac{e^{i\phi(N)} \det(e^{i\phi})}{\sqrt{\det C(\phi)}}, \quad (19)$$

$$y(\phi) = \frac{x(\phi)}{\int dg x(\phi)}, \quad \int dg y(\phi) = 1, \quad (20)$$

and

$$Y(\phi) = ie^{-i\phi} \sin \phi C(\phi) - i \int d\phi' y(\phi') e^{-i\phi'} \sin \phi' C(\phi'). \quad (21)$$

The quantities ρ and κ in the above equations are the HFB density-matrix and the pairing-tensor. e' in (8) are the single-particle energies of the cranked-deformed potential (1) and $\bar{v}$ in (12-14) is the uncoupled antisymmetric matrix-element of the two-body delta-interaction (5).

The term designated by Λ in (10) does not appear in the ordinary HFB formalism and it can be immediately shown that it vanishes for the gauge-angle $\phi = 0$. This term originates from the variation of the pairing-energy with respect to the density-matrix. In normal HFB, the pairing-energy depends only on the pairing-tensor, but the PHFB pairing-energy also depends on the density-matrix through the norm-overlap. Actually, in the general the norm-overlap depends on both density-matrix and the pairing-tensor [8]. But for the special case of number-projection, the term in the overlap-matrix which depends on the pairing-tensor can be rewritten in terms of the density-matrix by using the HFB relation $(\rho - \rho^2 = \kappa \kappa^\dagger)$.

Due to this transformation, the expression for Δ in (11) has a very simple appearance and reduces to the familiar form in the canonical representation [8,11].

The integration in (8-11) over the gauge-angle has been performed using the Gauss-Chebyshev quadrature method [12]. In this method, the integration over the gauge-angle is replaced by a summation. It can be shown [12] that the optimal number of mesh-points in the summation which eliminates all the components having undesired particle numbers is given by

$$M = \max\left(\frac{1}{2}N, \Omega - \frac{1}{2}\right) + 1, \quad (22)$$

where N is the number of particles and Ω is the degeneracy of the single-j shell. In the present study with $N=6$ and $\Omega = 6$, the optimal number of points required is $M = 4$.

In the present analysis of a single-j shell, the basis in which the HFB matrix is constructed are the magnetic sub-states of $j = 11/2$ with $m = (11/2, 9/2, \dots, -9/2, -11/2)$. The summation indices n_1, n_2, n_3 and n_4 in all the expressions given above run over these magnetic states. In order to check the dependence of the HFB and PHFB results on the pairing interaction, the calculations have been performed with monopole ($L=0$), monopole plus quadrupole ($L=0$ and 2), and with full delta-interaction. The results of the HFB and PHFB will be compared with the exact results for the three pairing-interactions.

The results of the cranking calculations with monopole-pairing force are compared in Fig. 1. In the three-panels of the figure, we compare the total-energy (E_{tot}), the pairing-energy (E_{pair}) and the alignment ($\langle J_x \rangle$) which is the expectation value of the angular-momentum along the rotational x-axis, as a function of the rotational frequency. The expressions for the total energy is given by

$$E_{tot} = \int d\phi y(\phi) (H_{sp}(\phi) + H_{ph}(\phi) + H_{pp}(\phi)), \quad (23)$$

where

$$H_{sp}(\phi) = \text{Tr} (e\rho(\phi)), \quad (24)$$

$$H_{ph}(\phi) = \frac{1}{2}\text{Tr} (\Gamma(\phi)\rho(\phi)), \quad (25)$$

$$H_{pp}(\phi) = -\frac{1}{2}\text{Tr}(\Delta(\phi)\bar{\kappa}^*(\phi)). \quad (26)$$

The expressions for the pairing-energy and the alignment are given by

$$E_{pair} = \int d\phi H_{pp}(\phi), \quad (27)$$

$$< J_x > = \int d\phi \text{Tr}(J_x \rho(\phi)). \quad (28)$$

It can be easily shown that for the gauge-angle, $\phi = 0$, the normal HFB expressions for these quantities are recovered.

It is apparent from the top pannel of Fig. 1 that the results of the exact shell model and the PHFB are very similar, the two curves are almost indistinguishable for all the frequency points. The results of HFB on the other hand deviate considerably from the exact results at lower frequencies. However, it can be seen from Fig. 2 that the HFB results converge towards the exact results with increasing rotational frequency. The HFB energy before the bandcrossing at $\hbar\omega = 0.5G$ is shifted from the exact energy by a constant amount and can be improved by renormalising the strength of the pairing-interaction. Therefore for the total energy, the HFB approach is not a poor approximation. The actual problem in HFB lies in the analysis of the pairing-energy and the alignment which are shown in the two lower-pannels of Fig. 1. The HFB pairing-energy has a finite value till $\hbar\omega = 0.45G$ and then suddenly goes to zero at $\hbar\omega = 0.5G$. This transition is an artefact of the HFB approach as is clearly evident from Fig. 1. The PHFB pairing-energy does drop at the bandcrossing but has a finite value at all the rotational frequencies. This feature of the pairing-energy is also reflected in the alignment, which is determined by the competition between the Coriolis forces and the pairing-correlations. The alignment till $\hbar\omega = 0.45G$ is very similar in all the three cases but then HFB value suddenly jumps to $< J_x > = 10$, the exact and the PHFB values on the other hand do not show this sharp transition. The gain in alignment at the first-crossing, referred to as the (AB)-crossing, is quite similar in all the cases, however, in HFB the crossing occurs somewhat earlier.

The results of pairing-interaction with monopole plus quadrupole terms are shown in Fig. 2. The comparison among HFB, PHFB and the exact results is quite similar to Fig. 2.

The total HFB energy is shifted by a constant factor from exact and PHFB energies before the bandcrossing. After the crossing, the HFB results become closer to the exact. The HFB pairing-energy depicts a transition and PHFB pairing-energy on the other hand drops smoothly at the bandcrossing. However, the HFB gain in alignment at the (AB)-crossing is much lower than 10 and has clearly a wrong behaviour after the crossing.

The results with the full delta-interaction are presented in Fig. 3. The HFB total-energy is now in a better agreement with exact and PHFB as compared to the results shown in Figs. 1 and 2. In fact, it is evident by comparing the three figures that the total-energy improves by including higher multipoles in the pairing-interaction, maximum deviation is noted for the monopole case. This appears to be in contradiction to our basic understanding of the mean-field approach, one normally expects that HFB or BCS is a better approximation for the pure monopole-pairing. However, it should be mentioned that in our HFB and PHFB analysis, we have included all the terms in the Hamiltonian. In particular, the particle-hole contribution (Γ) amounts to about 6G in the total-energy and is maximum with the full delta-interaction. If one excludes this contribution, the discrepancy would be largest for the delta-interaction.

The pairing-energy in Fig. 3 again depicts a phase transition at $\hbar\omega = 0.55G$ which is slightly higher than with monopole-interaction. The HFB (AB)-crossing with full delta-interaction is now close to the exact and PHFB. The gain in alignment at the (AB)-crossing is lower than 10 as in the case of monopole and quadrupole pairing force. The overall agreement with full delta-interaction appears to be better for HFB. The good agreement between PHFB and exact on the other hand is independent of the interaction used.

To conclude, in the present work the number-projected HFB approach recently developed has been applied to an exactly soluble cranked-deformed shell Hamiltonian. The main motivation has been to check the numerical applicability of the projection method. It is clear from the present study that the projected Hartree-Fock-Bogoliubov approach gives an accurate description of the yrast-states of the model Hamiltonian. The transition from a superfluid to the normal phase obtained in the HFB theory is shown to be smeared out with

the projection.

We would like to stress that the main advantage of the present projection method is that it has the same structure as that of normal HFB equations. Therefore, one can use the existing HFB computer codes and only the expressions for the Hartree-Fock potential and the pairing-field need to be redefined. Instead of the normal HFB fields, in the projection method one needs to calculate the projected quantities as given by (8-11). In the present model study, we find that the numerical work involved in the projection is similar to performing the bare HFB calculations. For each rotational frequency, the average CPU time on Pentium (166MHz) was 6.14s with projection as compared to 5.97s for normal HFB. The present projection method, therefore, preserves all the mathematical and computational simplicity of the HFB mean-field approach.

REFERENCES

- [1] Ring P. and Schuck P., *The Nuclear Many-Body Problem* (Springer Verlag, New York, 1980)
- [2] Magierski P., Cwiok S., Dobaczewski J., and Nazarewicz W., Phys. Rev. C 1993. V. 48. P. 1681.
- [3] Gall G., Bonche B., Dobaczewski J., Flocard H., Heenen P.-H., Z. Phys. A 1994 V. 348. P. 183.
- [4] Valor A., Egido J.L., and Robledo R., Phys. Rev. C 1996. V. 53. P. 172
- [5] Flocard H. and Onishi N., Ann. Phys. (N.Y.) 1996. V. 254 P. 275
- [6] Egido J.L. and Ring P., Nucl. Phys. A 1982. V. 383. P. 189
- [7] Egido J.L. and Ring P., Nucl. Phys. A 1982. V. 388. P. 19
- [8] Sheikh J.A. and Ring P., Nucl. Phys. A 2000 V. 665. P. 71.
- [9] Sheikh J.A., Nagarajan N.A., Rowley N. and Pal K.F., Phys. Lett. B 1989. V. 223. P. 1.
- [10] Brussaard J.P and Glaudemans P.W.M, *Shell Model Applications in Nuclear Spectroscopy* (North-Holland, Amsterdam, 1977)
- [11] Dietrich K., Mang H.J. and Pradal J.H., Phys. Rev. 1964. V. 135. P. B22
- [12] Hara K., Iwasaki S. and Tanabe K., Nucl. Phys. A 1979. V. 332. P. 69.

FIGURES

FIG. 1. The results of the total energy (E_{tot}), the pair-energy (E_{pair}) and the alignment (J_x) for six-particles in a deformed $j = 11/2$ orbital using the monopole-interaction. The PHFB results are indistinguishable from the exact shell models results.

FIG. 2. The results of the total energy (E_{tot}), the pair-energy (E_{pair}) and the alignment (J_x) for six-particles in a deformed $j = 11/2$ orbital using the monopole plus quadrupole interaction.

FIG. 3. The results of the total energy (E_{tot}), the pair-energy (E_{pair}) and the alignment (J_x) for six-particles in a deformed $j = 11/2$ orbital using the full delta-interaction.





